

PRVPATENT- OCH REGISTRERINGSVERKET
Patentavdelningen**Intyg
Certificate**

Härmed intygas att bifogade kopior överensstämmer med de handlingar som ursprungligen ingivits till Patent- och registreringsverket i nedannämnda ansökan.

Ansökan ingavs ursprungligen på engelska.

This is to certify that the annexed is a true copy of the documents as originally filed with the Patent- and Registration Office in connection with the following patent application.

The application was originally filed in English.

(71) Sökande AstraZeneca AB, Södertälje SE
Applicant (s)

(21) Patentansökningsnummer 0301119-4
Patent application number

(86) Ingivningsdatum 2003-04-14
Date of filing


Stockholm, 2004-04-26

För Patent- och registreringsverket
For the Patent- and Registration Office

Marita Öun

Avgift
Fee

PRIORITY DOCUMENT
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH
RULE 17.1(a) OR (b)



NEW NON-IONIC SURFACTANTS FOR SOLUBILIZING POORLY SOLUBLE MOLECULES.

Technical field

- 5 This invention relates to polyoxyalkylene glycols (in the following sometimes referred to as POAG) or monoalkylated polyoxyalkylene glycols esterified with O-acylated, O-alkylated or O-alkenylated hydroxy fatty acids, as well as their manufacture, their use in formulations (including pharmaceutical formulations) , and their use as surfactants.

10 Background of the invention

- The introduction of HTS methods (High Throughput Screening) in early drug discovery together with an enhanced demand on selectivity have in recent years increased the number of candidate drugs with a low aqueous solubility. In order to minimise administration volumes and obtain a high bioavailability it is of great practical importance for
15 pharmaceutical formulators to have the ability to increase the solubility of these compounds when suitable dosage forms are developed. These preparations can be intended both for assessment of medical effect in humans and safety studies in animals during the development of a new drug, and as the final pharmaceutical dosage form for the marketed product.

20

One commonly used method to increase the solubility of poorly soluble compounds is to solubilise the compound in a micellar system by the use of surfactants. ["The Theory and Practice of Industrial Pharmacy" 2nd ed. Lea & Febiger, 1976, p. 108-111.]

- 25 The main advantages with micellar systems are the stability over a wide composition range, simplicity of preparation, low viscosity and the fact that a micellar system is a thermodynamically stable single phase which is optically clear. Surfactants can be divided into anionic, e.g. sodium lauryl sulfate, cationic, e.g. cetyl trimethyl-ammonium bromide, zwitter-ionic, e.g. alkyl betaines and non-ionic surfactants, according to their chemical
30 properties.

The choice of surfactants for use in pharmaceutical applications depends to some extent on the route of administration and is rather limited since most surface-active compounds are not tolerated well enough for pharmaceutical use. For parenteral use ionic surfactants are not suitable since these cause hemolysis of red blood cells and destruction of T lymphocyte cells at low concentrations. [“Solubility & Solubilization in aqueous media.”, Yalkowsky, 1999]. The most accepted surfactants for parenteral use are phospholipids and non-ionic surfactants. For oral use non-ionic surfactants are usually preferred but ionic-surfactants have been used in low concentrations.

Non-ionic surfactants used in pharmaceutical applications today include substances/mixtures such as ethoxylated castor oil (Cremophor EL), ethoxylated sorbitan fatty acid esters, e.g. polyoxyethylene sorbitan monooleate (Tween 80), sorbitan fatty acid esters, e.g. sorbitan monooleate (Span 80), ethoxylated hydroxystearic acid, e.g. polyethylene glycol 660 (12-)hydroxystearate (Solutol HS15), ethylene and propylene oxide block copolymers (Pluronic F68) and fatty acid esters of glycerol (Imwitor 742).

The above described non-ionic surfactants which are presently used in pharmaceutical applications do, however, exhibit a number of disadvantages.

For example, the commercial non-ionic surfactants available for pharmaceutical formulators are complex mixtures of different molecules which makes the characterisation of these products very difficult, giving an expensive and tedious analytical process to ensure adequate quality (for i.a. pharmaceutical applications).

Recent studies on adverse effects on epithelial cells have shown that commercial non-ionic surfactants have a profound effect on epithelial cells in concentrations typically used for solubilisation (Östh, Karin, Thesis: The horizontal Ussing chamber method in studies of nasal drug delivery, 2002. Faculty of Pharmacy, Uppsala University).

It is also well known that surface-active compounds often cause hemolysis at low concentrations when administered parenterally.

The existing non-ionic surfactants systems used for parenteral administration are all based on polyethylene glycol derivatives. Although there are several pharmaceutical products for parenteral administration on the market containing these surfactants, they all suffer from quite severe side effects, like release of histamine which in severe cases can lead to anaphylactic shocks (Lorentz et al., Agents and Actions, Vol 12, 1/2, 1982). anaphylactic shocks).

Histamine release is believed to be caused by impurities in the commercial products and since the non-ionic surfactants used are very complex mixtures of different molecules it is not possible to purify existing products. Also in such situations it is hard to relate any side-effects to a particular molecule. (Vulfsson. In "Novel Surfactants". Holmberg editor. Marcel Dekker 1988. p. 279-97.)

In EP 0017059 A1, reaction products of monohydroxy fatty acids with ethylene oxide in a given molar ratio are mentioned (the Solutol® type of compounds).

The products formed are mixtures of monoesters or diesters of polyethylene glycol (PEG) and monohydroxy fatty acids or estolides, the latter commonly known as a generic name for linear oligomeric polyesters of hydroxyl fatty acids wherein the carboxyl group and hydroxyl group of hydroxyl fatty acids are dehydrated to form oligomers. The products of EP 0017059A1 comprising two or more monohydroxy fatty acids where the monohydroxy fatty acid of the estolide may either be attached directly to the hydroxyl group of another fatty acid or to a hydroxyl group of a PEG chain attached to the aforementioned hydroxyl group of a monohydroxy fatty acid. These reaction products are stated to be used especially as dissolution enhancers for pharmaceutical purpose. With these kind of compounds, the resulting synthesis product will always be a mixture of compounds. Solutol HS 15 is such a product. Short PEG chains (one type of polyoxyalkylene glycol, or POAG, chain) are a characteristic feature of the compounds claimed in EP 0017059 A1.

US 6,365,637 claims the use of esters or amides of hydroxylated carboxylic acids as solubilizers, for i.a. pharmaceutical purposes. These compounds all have short PEG chains. Furthermore, the optional use of a dimerized fatty acid, as described in US 6,365,637, of

commercial quality including both monomeric, dimeric, trimeric and higher polymerized acids in the synthesis is a draw-back when one desires to obtain highly pure compounds.

5 Description of the invention

Accordingly, there is a need for new effective non-ionic surface-active compounds not having the above mentioned disadvantages of being complex mixtures, or potential for inducing adverse reactions seen as e.g. epithelial cell interaction, histamine release or
10 hemolysis.

As some side-effects are believed to be caused by impurities in the commercial products and as the non-ionic surfactants used are very complex mixtures of different molecules it is not possible to purify existing products (Vulfsson. In "Novel Surfactants". Holmberg
15 editor. Marcel Dekker 1988. p. 279-97.). Therefore, there is need for new "non-toxic" and well-defined surfactants having high solubilization capacity, for use as solubilizers of poorly soluble drug molecules. Such compounds may have use in both pharmaceutical and other fields.

20 It is also a need for a method of synthesis that permits the surfactant to be manufactured as a highly pure compound.

Furthermore, there is also a need for these compounds with an improved profile regarding side-effects and having high solubilization capacity as surfactants, for use in formulations
25 generally, but especially for use in pharmaceutical formulations.

It has now surprisingly been found that compounds being an O-acylated or O-alkylated/ O-alkenylated hydroxy-fatty acid (hydroxy-fatty acid in the following also referred to as "HFA"), esterified with polyoxyalkylene glycol (POAG) or monoalkylated POAG, where
30 the POAG (or its derivative) has a specified average chain-length are better tolerated than compounds of the prior art, especially with regard to low haemolytic activity and lack of interaction with epithelial cells (CACO-2 cells). Furthermore they can be prepared as well-

defined compounds and they are very effective as solubilisers. Such compounds can advantageously be used in i.a. pharmaceutical formulations such as tablets, capsules, powders, dispersions, emulsions, rectal formulations like suppositories and also in solutions. In addition they can be used as e.g. media for making solid dispersions, absorption enhancers, emulsifiers in emulsions and microemulsions, dispersing agents in solid dispersions, emulsifier in self- emulsifying systems, lubricants in tablet pressing, wetting agents in granulation processes, vehicles in spray-drying compositions etc, are also uses, not limited to any particular administration route, that are contemplated in this invention.

10


It has also been found that a process involving an enzymatic step connecting a previously formed O-acyl HFA or O-alkyl/ O-alkenyl HFA (or esters of such a HFA or their derivatives) with POAG or alkylPOAG (e.g. MePEG1200, also known as polyethylene glycol monomethyl ether with average molecular weight 1200) using a hydrolytic enzyme, said enzyme having the capability of catalyzing ester formation between the carboxylic group of the HFA-derivative and the ending hydroxyl group of POAG or POAG-derivative, without catalyzing any reaction with existing ester or ether bond on the O-acyl/alkyl/alkenyl-HFA (or corresponding derivative). As an example, lipase B from *Candida antarctica* or an equivalent can be used. This enzyme is particularly beneficial to employ to obtain more homogenous reaction products.

20

Detailed description of the invention

The compounds of the invention, surprisingly having the beneficial advantages of better tolerability, low hemolytic activity, possibility to be produced with higher purity and having high solubilization capacity, have chemical structures as described in formula (I) below;

30



(I)

The compounds are based on an acylated/alkylated/alkenylated hydroxy fatty acid forming an ester bond with a poly (oxyalkylene) glycol [in a shorter form POAG] or monoalkylated poly (oxyalkylene) glycol on the carboxylic group end.

5

The R_1 group positioned on the outer end of the POAG chain may be H or $C_1 - C_4$ alkyl. In one aspect of the invention R_1 is H, alternatively R_1 is $C_1 - C_4$ alkyl. In one preferred embodiment of the invention, the R_1 is H or $C_1 - C_2$ alkyl. In a more preferred embodiment of the invention R_1 is $C_1 - C_2$ alkyl. In a most preferred embodiment the R_1 is methyl.

10

The term POAG includes poly (oxyethylene) glycols, commonly known as i.a. PEG's, and poly (oxypropylene) glycols, commonly known as PPG's. The oxypropylene units may be linear or branched.

15 Thus, R_3 is chosen among ethylene, propylene or branched propylene.

The POAG chain used in the invention has preferably, counted as poly (ethylene) glycol, an average molecular weight of 1100 – 20000. More preferred is to use a POAG with an average molecular weight of 1100 – 10000, most preferred is to use a POAG with an average molecular weight of 1100 – 2500, all these average molecular weights counted as for poly (ethylene) glycol.

20

These average molecular weights are counted as excluding the weight of any alkyl groups in the alkyl-derivatized POAG's.

25

The poly (oxyalkylene) glycols or monoalkylated poly (oxyalkylene) glycols used are of pure, monodisperse or polydisperse commercial quality.

30

The above range limits given for polyoxyalkylene glycol (e.g. PEG) chain average molecular weight, corresponds to values of z being 25 – 455, preferably 25 – 228 and most preferably 25-57.

Table 1 below shows the relation between z and average molecular weight for PEG and PPG, respectively.

Table 1.

z	Molecular weight Polyoxyethylene Glycol (PEG)	Molecular weight Polyoxypropylene Glycol (PPG)	Remark
1	44	58	As monomer unit
25	1100	1450	As multiples of monomer units
57	2508	3306	"
100	4400	5800	"
228	10032	13224	"
455	20020	26390	"
500	22000	29000	"

5

The carbon chain in the hydroxy fatty acid part of the inventive compounds are characterized in that the oxygen from the hydroxy group is positioned such as that x is 2 - 18 and y is 1 to 17, while the sum of $(x + y)$, defining the length of the hydroxy fatty acid carbon chain, is 3 - 19.

10

More preferably the value of x is 2 - 15 and the value of y is 4 - 17, while the sum of $(x + y)$ is 6 - 19 .

15

Most preferably the value of x is 2 - 12 and the value of y is 7 - 17, while the sum of $(x + y)$ is 9 - 19 .

20

The oxygen in the hydroxyl group on the HFA residue is connected (via an ester linkage or ether linkage) to a group R_2 , R_2 being C_{14} to C_{22} , linear or branched, acyl, alkyl or alkenyl wherein the acyl, alkyl or alkenyl may be optionally further substituted with one or more of the following (independently selected); halogen, cyano, carboxy, carbamoyl,

carbamoyl(C₁-C₄)alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, mercapto, nitro, amino, (C₁-C₄)alkylamino, phenyl, naphthyl, phenyloxy, naphthyloxy, (C₁-C₄)alkylthio, or (C₁-C₄)alkylsulfinyl.

- 5 In one embodiment R₂ is unsubstituted. In another embodiment R₂ is substituted by one or two substituents, preferably one.

In another second embodiment of the invention, the R₁ group positioned on the outer end of the POAG chain is C₁ - C₄ alkyl. In one more preferred second embodiment of the invention, R₁ is C₁ - C₂ alkyl. In the especially most preferred second embodiment R₁ is methyl.

The term POAG includes poly (oxyethylene) glycols and poly (oxypropylene) glycols. The oxypropylene units may be linear or branched.

15

Thus, R₃ is chosen among ethylene, propylene or branched propylene.

The POAG chain used in the invention has preferably, counted as poly (ethylene) glycol, an average molecular weight of 1100 - 20000. More preferred is to use a POAG with an average molecular weight of 1100 - 10000, most preferred is to use a POAG with an average molecular weight of 1100 - 2500, all these average molecular weights counted as for poly (ethylene) glycol.

20

These average molecular weights are counted as excluding the weight of any alkyl groups in the alkyl-derivatized POAG's.

25

The poly (oxyalkylene) glycols or monoalkylated poly (oxyalkylene) glycols used are of pure, monodisperse or polydisperse commercial quality.

30

The above range limits given for polyoxyalkylene glycol (e.g. PEG) chain average molecular weight, corresponds to values of z being preferably 25 - 455, more preferably 25 - 228 and most preferably 25-57.

Table 1 above shows the relation between z and average molecular weight for PEG and PPG, respectively.

- 5 The carbon chain in the hydroxy fatty acid part of the inventive compounds are characterized in that the oxygen from the hydroxy group is positioned such as that x is 2 - 18 and y is 1 to 17, while the sum of $(x + y)$, defining the length of the hydroxy fatty acid carbon chain, is 3 - 19.
- 10 More preferably the value of x is 2 - 15 and the value of y is 4 - 17, while the sum of $(x + y)$ is 6 - 19 .


Most preferably the value of x is 2 - 12 and the value of y is 7 - 17, while the sum of $(x + y)$ is 9 - 19 .

15

- The oxygen in the hydroxyl group on the HFA residue is connected (via an ester linkage or ether linkage) to a group R_2 , R_2 being C_{14} to C_{22} , linear or branched, acyl, alkyl or alkenyl wherein the acyl, alkyl or alkenyl may be optionally further substituted with one or more of the following; halogen, cyano, carboxy, carbamoyl, carbamoyl(C_1 - C_4)alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, mercapto, nitro, amino, (C_1 - C_4)alkylamino, phenyl, naphthyl, phenyloxy, naphthyloxy, (C_1 - C_4)alkylthio, or (C_1 - C_4)alkylsulfinyl.
- 20

In one second embodiment R_2 is unsubstituted. In another second embodiment R_2 is substituted by one or two substituents, preferably one.

25



Synthesis

Ester derivatization of hydroxy fatty acid

The synthesis can, for example, be performed by starting from the hydroxy fatty acid or its
5 C₁ – C₄ alkyl esters. If needed, the hydroxy fatty acid starting material may be purified by a
suitable method, e.g. extraction or chromatographic methods. This is beneficial for
obtaining the best results of the invention.

10 Esterification of the fatty acid hydroxyl group can for example be performed with an acyl
chloride. The two reactants are mixed in a suitable solvent, e.g. methyl tert-butyl ether
(MTBE) containing pyridine in a suitable concentration e.g. 1.5 equivalents, in relation to
the acyl chloride.

Purification of this product can for example be done by extraction. Suitably, the product is
15 first washed with a weakly acidic solution, e.g. 1% sulphuric acid, and then with a weakly
basic solution, e.g. saturated aqueous sodium bicarbonate. If needed, further purification
may be done using a chromatographic method, e.g. preparative silica gel chromatography.

20 Other methods known in the state of the art can equally well be used to accomplish
esterification of the fatty acid hydroxyl group.

Ether derivatization of hydroxy fatty acid

The synthesis can, for example, be performed by starting from esters of the hydroxy fatty
acid. If needed, the hydroxy fatty acid ester may be purified by a suitable method, e.g.
25 extraction or chromatographic methods. This is beneficial for obtaining the best results of
the invention.

30 Etherification of the fatty acid hydroxyl group with an alcohol can for example be
performed in two steps. The first step may consist of reacting the alcohol with toluene-4-
sulfonyl chloride in a suitable solvent, e.g. dichloromethane, containing pyridine in a
suitable concentration e.g. 1.5 equivalents in relation to the toluene-4-sulfonyl chloride.

Purification of the produced alcohol toluene-4-sulfonate can for example be done by

extraction. Suitably, the product is then washed with a weakly acidic solution, e.g. 1% sulphuric acid. If needed, further purification may be done using a chromatographic method, e.g. preparative silica gel chromatography.

- 5 The second step may consist of reacting the alcohol toluene-4-sulfonate with the hydroxyl fatty acid ester in a suitable system of solvent, base and catalyst, e.g. in acetonitrile containing 1 equivalent potassium carbonate and 0.05 equivalents sodium iodide, in relation to the alcohol toluene-4-sulfonate. Purification of the produced alkyloxy acid ester can for example be done by extraction. Suitably, the product is then washed with a weakly
10 acidic solution, e.g. 1% sulphuric acid. If needed, further purification may be done using a chromatographic method, e.g. preparative silica gel chromatography.

Other methods known in the state of the art can equally well be used to accomplish etherification of the fatty acid hydroxyl group.

15

Esterification with POAG or POAG derivatives

One very advantageous succeeding step after the esterification or etherification of the HFA, yielding surprisingly pure products at high yield is when the obtained HFA derivatives are esterified with the polyoxyalkylene glycol, or monoalkylated

- 20 polyoxyalkylene glycol using a hydrolytic enzyme, and this enzyme is having the capability of catalyzing ester formation between the carboxylic group of the HFA derivative and the ending hydroxyl group of POAG or POAG-derivative, without catalyzing any reaction with existing ester or ether bond on the O-acyl/alkyl/alkenyl-HFA. This is in the following some times referred shortly to as "enzymatic POAGylation". As an
25 example, lipase B from *Candida antarctica* or an equivalent can be used. A preferred enzyme to be used in the inventive process is lipase B from *Candida antarctica*. The most preferred form of the enzyme to be used in the inventive process is the immobilized form of the lipase B enzyme from *Candida antarctica*.

- 30 The enzymatic POAGylation step is advantageously performed in combination with the use of vacuum to remove water or any other volatile co-product formed during the esterification.

The reaction can be monitored by using HPLC. A preferred option is to choose reversed phase chromatography, using an evaporative light scattering detector.

- 5 Reaction temperatures when utilizing lipase B from *Candida antarctica*, are typically between 50 and 90 degrees Celsius.

Formulations

10

The compounds of the invention may be incorporated for different purposes in formulations, thereby taking advantage of their benefits as described above.

Such purposes include usages as; surface active agents, solubilizers, detergents, dispersants for liquid dispersions, solid dispersants, emulsifiers, carriers, freeze-drying additives,

- 15 spray-drying additives and wetting agents.

The compounds of the invention can be used in combination with any other compound not being negatively influenced by their presence. Especially it is foreseen to use them in combination with compounds in water being (according to definition given in US

- 20 Pharmacopoeia 24, from 2000, page 10) sparingly soluble or less soluble. This includes sparingly soluble, slightly soluble, very slightly soluble, practically insoluble and insoluble, according to the same definition. Thus it includes compounds having a solubility of less than approx. 0.033 mg/mg in a solvent, corresponding to 33 mg/ml in water. The solvent in relation to this invention is water, at 25 degrees Celsius and normal atmospheric pressure.
- 25

Pharmaceutical formulations

- 30 The compounds of the invention may be incorporated in pharmaceutical formulations for any purpose known according to the art, thereby taking advantage of their benefits as described above. Such purposes include, but are not limited to using them as; surface

active agents, solubilizers, emulsifiers, carriers, solid dispersants, dispersants for liquid dispersions, wetting agents, lubricants, freeze-drying additives, spray-drying additives, etc.

As such they may be included in parenteral formulations (e.g. subcutaneous, intravenous, intramuscular, intraperitoneal, intraarterial, intracerebrovascular), oral formulations, rectal formulations, topical formulations, buccal formulations, sublingual formulations, this enumeration not intended to be limiting in any way.

Preferred formulations are oral, topical, rectal or parenteral. More preferred formulations are oral or parenteral. Most preferred is parenteral formulations.

Non-limiting examples of oral formulations suitable for use with compounds of the invention are tablets, effervescent tablets, capsules, granules, pellets, powders, sachets, dispersions, suspensions, emulsions, microemulsions, self-emulsifying systems and solutions.

In the use of the compounds according to the invention, the formulator may need to include various excipients, such as cosolvents, buffers, polymers, disintegrants, fillers/diluents, stabilisers and preservatives.

20

The compounds of the invention can be used in combination with any pharmaceutically active ingredient (drug) or vitamin not being negatively influenced by their presence. Especially is it foreseen to use them in combination with drugs or vitamins in water being (according to definition given in US Pharmacopoeia 24, from 2000, page 10) sparingly soluble or less soluble. This includes sparingly soluble, slightly soluble, very slightly soluble, practically insoluble and insoluble, according to the same definition. Thus it includes drugs or vitamins having a solubility of less than approx. 33 mg/mg in a solvent, corresponding to 33 mg/ml in water. The solvent in relation to this invention is water, at 25 degrees Celsius and normal atmospheric pressure.

30

According to a further embodiment the pharmaceutical formulations may comprise one or more compound according to the invention, and may also contain more than one

pharmaceutically active ingredient. It is further possible for the pharmaceutical formulations (utilizing the invention) to comprise one or more vitamin. Another aspect that is contemplated is that combinations of one or more vitamin with one or more pharmaceutically active ingredient in may be used in the formulations of the invention.

5

A preferred embodiment of pharmaceutical formulation according to the invention comprises one or more drug or pharmaceutically active ingredient selected among; Proton pump inhibitors, calcium channel blockers, adrenergic beta-blockers, anesthetics, steroids, antioxidants, renin inhibitors, alkaloids, cytostatics, anticoagulants, lipid
10 regulating agents, anti-depressants, neuroleptics, immunosuppressants, immunomodulators, antibiotics and non-steroidal anti-inflammatory agents.

The following examples are provided to illustrate the present invention.

15

Example 1.

Synthesis of PEG600 mono-12-lauroxyloxy-stearate.

20 In general the reaction followed the way ;

12-hydroxystearic acid → ethyl 12-hydroxystearate → ethyl 12-lauroxyloxy-stearate →
PEG600 mono-12-lauryloxy-stearate.

Ethyl 12-hydroxystearate.

25 Technical grade 12-hydroxystearic acid (purchased from Aldrich) was purified to >99% by extraction with hexane.

150 mmol (45 g) of the purified 12-hydroxystearic acid and 5 mmol (0.5 g) sulfuric acid was dissolved in 300 ml ethanol in a 500 ml round flask with a Dimroth condenser and
30 refluxed at 78 °C for 20 hours. The product was isolated by solvent evaporation and subsequent dissolution in 300 ml methyl tert-butyl ether (hereinafter MTBE). The liquid

was extracted with water (3x100 ml) and then evaporated. The product was recovered in 96% isolated yield (47.8 g).

Ethyl 12-lauroyloxy-stearate.

- 5 20 mmol (6.57 g) of the obtained ethyl 12-hydroxystearate and 30 mmol (2.4 g) pyridine was dissolved in 200 ml MTBE at 50 °C in a 250 ml round flask with a Dimroth condenser and drying tube. Under stirring, 19.4 mmol (4.24 g) lauroyl chloride (from Sigma Aldrich) was added slowly and allowed to react for 5 hours. The reaction was stopped by lowering the temperature to 25 °C and adding 100 ml water.
- 10 The solution was extracted three times with 50 ml portions of 1% aqueous sulfuric acid, followed by extraction with saturated aqueous sodium bicarbonate (90 g/l, 3x50 ml) and finally water (3x50 ml). Remaining solvent was removed by evaporation. Released lauric acid was removed by chromatography on silica gel eluting with 5% ethyl acetate in cyclohexane with 0.1% acetic acid. Afterwards the solvent was evaporated.
- 15 Ethyl 12-lauroyloxy-stearate was recovered in 71% (7.1 g) isolated yield.

PEG600 mono-12-lauroyloxy-stearate.

- 7.5 mmol (3.7 g) of the obtained ethyl 12-lauroyloxy-stearate and 75 mmol (45 g) PEG600 were mixed with 50 mg immobilized lipase B from *C. antarctica* (Novozym 435 from
- 20 Novozymes A/S Denmark) in a 250 ml round flask with stirring under vacuum (<1 mm Hg) at 60 °C for 16 h.

- The reaction was monitored using reversed phase HPLC with a Supelco Discovery C18-column (4.6 mm i.d. x 150 mm), 1 ml/min of a 95:5 methanol:water mixture with addition
- 25 of 0.5% acetic acid in the separation. Detection was done with a Sedex 45 evaporative light scattering detector (Sedere, Alfortville, France) at 30 °C and 2 bar air pressure.

- Afterwards the enzyme was removed by filtration and the product was extracted with ethyl acetate and saturated aqueous sodium chloride according to ISO-2268 ["Surface active
- 30 agents (non-ionic) - Determination of polyethylene glycols and non-ionic active matter (adducts) - Weibull method", International Standard Organization, ISO 2268:1972] but at room temperature followed by solvent evaporation, dissolution in dry ethyl acetate,

filtration and evaporation. PEG600 mono-12-lauroxyloxy-stearate was recovered in 93% (7.2 g) isolated yield.

5 Example 2

Synthesis of PEG600 mono-12-propionyloxy-stearate.

In general the reaction followed the way ;

12-hydroxystearic acid → 12-propionyloxy-stearic acid → PEG600 mono-12-propionyloxy-
10 stearate.

12-Propionyloxy-stearic acid.

Technical grade 12-hydroxystearic acid (purchased from Aldrich) was purified to >99%
15 by extraction with hexane.

10 mmol (3.0 g) of the purified 12-hydroxystearic acid and 25 mmol (2.0 g) pyridine were dissolved in 150 ml MTBE at 50 °C in a 250 ml round flask with a Dimroth condenser and drying tube. Under stirring, 13 mmol propionyl chloride (from Sigma Aldrich) was added
20 slowly and allowed to react for 19 hours. The reaction was stopped by lowering the temperature to 25 °C and adding 50 ml water.

The solution was extracted with 1% aqueous sulfuric acid (3x25 ml) followed by saturated aqueous sodium bicarbonate (90 g/l, 3x25 ml) and water (3x25 ml). Remaining solvent was removed by evaporation. 12-Propionyloxy-stearic acid was recovered in 97% (3.47 g)
25 isolated yield.

PEG600 mono-12-propionyloxy-stearate.

9.9 mmol (3.7 g) 12-propionyloxy-stearic acid and 99 mmol (59.5 g) PEG600 were mixed with 50 mg immobilized lipase B from *C. antarctica* (Novozym 435 from Novozymes A/S
30 Denmark) in a 250 ml round flask with stirring under vacuum (<1 mm Hg) at 60 °C for 21 hours.

The enzyme was removed by filtration and the product was extracted with ethyl acetate and saturated aqueous sodium chloride according to ISO-2268 [1972] but at room temperature followed by solvent evaporation, dissolution in dry ethyl acetate, filtration and evaporation. PEG600 mono-12-propionyloxy-stearate was recovered in 78% (7.5 g) isolated yield.

Example 3

Synthesis of PEG1500 mono-12-stearoyloxy-stearate.

10

In general the reaction followed the way ;

12-hydroxystearic acid → ethyl 12-hydroxystearate → ethyl 12-stearoyloxy-stearate → PEG1500 mono-12-stearoyloxy-stearate.

15 The 12-hydroxystearic acid was from the same origin and purified as in the preceding examples.

Ethyl 12-hydroxystearate

Was obtained according to Example 1.

20

Ethyl 12-stearoyloxy-stearate

10 mmol (3.29 g) ethyl 12-hydroxystearate and 15 mmol (1.2 g) pyridine were dissolved in 100 ml MTBE at 50 °C in a 250 ml round flask with Dimroth condensers and drying tubes.

Under stirring, 9.7 mmol stearoyl chloride (2.94 g) was added slowly. The reaction was stopped after 15 hours by lowering the temperature to 25 °C and adding 50 ml water.

The solution was extracted with 1% aqueous sulfuric acid (3x25 ml) and remaining solvent was removed by evaporation.

Released stearic acid was removed by chromatography on silica gel eluting with a gradient from 5% to 10% ethyl acetate in cyclohexane fortified with 0.1% acetic acid followed by evaporation of the solvent.

Ethyl 12-stearoyloxy-stearate was recovered in 81% (4.7 g) isolated yield.

PEG1500 mono-12-stearoyloxy-stearate

7.8 mmol (4.7 g) ethyl 12-stearoyloxy-stearate was mixed with 100 mmol (150 g) PEG1500 and 50 mg immobilized *C. antarctica* lipase B in a 500 ml round flask with stirring under vacuum (<1 mm Hg) at 75 °C for 39 hours.

- 5 The enzyme was removed by filtration and the product was extracted with ethyl acetate and saturated aqueous sodium chloride according to ISO-2268 [1972] at room temperature followed by solvent evaporation, dissolution in dry ethyl acetate, filtration and evaporation. PEG1500 mono-12-stearoyloxy-stearate was recovered in 71% (11.4 g) isolated yield.

10

Example 4***Synthesis of PEG1500 mono-12-oleoyloxy-stearate.***

In general the reaction followed the way ;

- 15 12-hydroxystearic acid → ethyl 12-hydroxistearate → ethyl 12-oleoyloxy-stearate → PEG1500 mono-12-oleoyloxy-stearate.

- 20 The 12-hydroxystearic acid was from the same origin and purified as in the preceding examples.

Ethyl 12-hydroxystearate

Was obtained according to Example 1.

25 **Ethyl 12-oleoyloxy-stearate**

10 mmol (3.29 g) ethyl 12-hydroxystearate and 15 mmol (1.2 g) pyridine were dissolved in 100 ml MTBE at 50 °C in a 250 ml round flask with Dimroth condensers and drying tubes.

Under stirring, 9.7 mmol oleoyl chloride (2.92 g) was added slowly. The reaction was stopped after 15 hours by lowering the temperature to 25 °C and adding 50 ml water.

- 30 The solution was extracted with 1% aqueous sulfuric acid (3x25 ml) and remaining solvent was removed by evaporation.

Released oleic acid was removed by chromatography on silica gel eluting with a gradient from 5% to 10% ethyl acetate in cyclohexane fortified with 0.1% acetic acid followed by evaporation of the solvent.

Ethyl 12-oleoyloxy-stearate was recovered in 84% (4.9 g) isolated yield.

5

PEG1500 mono-12-oleoyloxy-stearate

8.2 mmol (4.9 g) ethyl 12-oleoyloxy-stearate was mixed with 100 mmol (150 g) PEG1500 and 50 mg immobilized *C. antarctica* lipase B in a 500 ml round flask with stirring under vacuum (<1 mm Hg) at 75 °C for 39 hours.

- 10 The enzyme was removed by filtration and the product was extracted with ethyl acetate and saturated aqueous sodium chloride according to ISO-2268 [1972] at room temperature followed by solvent evaporation, dissolution in dry ethyl acetate, filtration and evaporation. PEG1500 mono-12-oleoyloxy-stearate was recovered in 88% (14.8 g) isolated yield.

15

Example 5

Synthesis of MePEG1200 12-palmitoyloxy-stearate

In general the reaction followed the way ;

- 20 12-hydroxystearic acid → ethyl 12-hydroxystearate → ethyl 12-palmitoyloxy-stearate → MePEG1200 12-palmitoyloxy-stearate.

Ethyl 12-hydroxystearate

Was obtained according to Example 1.

25

Ethyl 12-palmitoyloxy-stearate

10 mmol (3.28 g) ethyl 12-hydroxystearate and 15 mmol (1.2 g) pyridine was dissolved in 100 ml MTBE at 50 °C in a 250 ml round flask with a Dimroth condenser and drying tube. Under stirring, 9.7 mmol (2.67 g) palmitoyl chloride was added slowly and allowed to react for 16 hours. The reaction was stopped by lowering the temperature to 25 °C and adding 50 ml water.

30

The solution was extracted with 1% aqueous sulfuric acid (3x25 ml) followed by saturated aqueous sodium bicarbonate (90 g/l, 3x25 ml) and water (3x25 ml). Remaining solvent was removed by evaporation.

- Released palmitic acid was removed by chromatography on silica gel eluting with 5% ethyl acetate in cyclohexane fortified with 0.1% acetic acid followed by evaporation of the solvent. Ethyl 12-palmitoyloxy-stearate was recovered in 87% (4.6 g) isolated yield.

MePEG1200 12-palmitoyloxy-stearate

- A portion of 3.76 g (6.97 mmol) ethyl 12-palmitoyloxy-stearate was mixed with 7.7 mmol MePEG1200 (9.2 g) and 50 mg Novozym 435 in a 100 ml round flask under vacuum (<1 mm Hg) with stirring at 75 °C for 200 hours.

The enzyme was removed by filtration and the product was extracted with ethyl acetate and saturated aqueous sodium chloride according to ISO-2268 [1972] at room temperature followed by solvent evaporation, dissolution in dry ethyl acetate, filtration and evaporation.

- MePEG1200 12-palmitoyloxy-stearate was recovered in 81% (9.75 g) isolated yield

Example 6

Synthesis of MePEG2000 12-palmitoyloxy-stearate

20

In general the reaction followed the way ;

12-hydroxystearic acid → ethyl 12-hydroxystearate → ethyl 12-palmitoyloxy-stearate → MePEG2000 12-palmitoyloxy-stearate.

25

Ethyl 12-hydroxystearate

Was obtained according to Example 1.

Ethyl 12-palmitoyloxy-stearate

30

10 mmol (3.28 g) ethyl 12-hydroxystearate and 15 mmol (1.2 g) pyridine was dissolved in 100 ml MTBE at 50 °C in a 250 ml round flask with a Dimroth condenser and drying tube. Under stirring, 9.7 mmol (2.67 g) palmitoyl chloride was added slowly and allowed to

react for 16 hours. The reaction was stopped by lowering the temperature to 25 °C and adding 50 ml water.

The solution was extracted with 1% aqueous sulfuric acid (3x25 ml) followed by saturated aqueous sodium bicarbonate (90 g/l, 3x25 ml) and water (3x25 ml). Remaining solvent

5 was removed by evaporation.

Released palmitic acid was removed by chromatography on silica gel eluting with 5% ethyl acetate in cyclohexane fortified with 0.1% acetic acid followed by evaporation of the solvent. Ethyl 12-palmitoyloxy-stearate was recovered in 87% (4.6 g) isolated yield.

10 MePEG2000 12-palmitoyloxy-stearate

A portion of 3.76 g (6.97 mmol) ethyl 12-palmitoyloxy-stearate was mixed with 7.7 mmol MePEG2000 (15.3 g) and 50 mg Novozym 435 in a 100 ml round flask under vacuum (<1 mm Hg) with stirring at 75 °C for 200 hours.

15 The enzyme was removed by filtration and the product was extracted with ethyl acetate and saturated aqueous sodium chloride according to ISO-2268 [1972] at room temperature followed by solvent evaporation, dissolution in dry ethyl acetate, filtration and evaporation. MePEG2000 12-palmitoyloxy-stearate was recovered in 80% (14.0 g) isolated yield.

20 **Example 7**

Synthesis of PEG1500 mono-12-palmitoyloxy-stearate

The general reaction route is: palmityl alcohol → palmityl toluene-4-sulfonate → ethyl 12-palmitoyloxy-stearate → PEG1500 mono-12-palmitoyloxy-stearate

25

Palmityl toluene-4-sulfonate

10 mmol palmityl alcohol and 20 mmol pyridine are dissolved in 100 ml dry dichloromethane. 10 mmol toluene-4-sulphonyl chloride is slowly added and the mixture is heated to 40 °C with stirring for 1 hour. The reaction is cooled to room temperature and 25 ml cold water is added to stop the reaction. The mixture is washed with 3x50 ml cold, 1% aqueous sulfuric acid to remove the pyridine followed by 3x25 ml cold water and evaporation to yield palmityl toluene-4-sulfonate.

30

Ethyl 12-hydroxystearate

Is obtained in accordance with Example 1.

5 Ethyl 12-palmitoxy-stearate

Dissolve 10 mmol palmityl toluene-4-sulfonate, 10 mmol ethyl 12-hydroxystearate, 10 mmol anhydrous potassium carbonate and 0.5 mmol sodium iodide in 100 ml dry acetonitrile and heat to 81 °C with stirring for 6 hours. The reaction is cooled to room temperature. The solvent is evaporated and the products are suspended in MTBE and

10 washed with 3x50 ml cold water. The solvent is evaporated to yield ethyl 12-palmitoxy-stearate.

PEG1500 mono-(12-palmitoxy-stearate)

10 mmol ethyl 12-palmitoxy-stearate is mixed with 100 mmol PEG1500 and 100 mg
15 immobilized *Candida antarctica* lipase B in a 500 ml round flask with stirring under vacuum (<1 mm Hg) at 75 °C for 40 hours.

The enzyme is removed by filtration and the product is extracted with ethyl acetate and saturated aqueous sodium chloride according to ISO-2268 [1972] followed by solvent evaporation, dissolution in dry ethyl acetate and filtration. The solvent is evaporated to
20 yield PEG1500 mono-(12-palmitoxy-stearate).

Example 8.***Synthesis of BuPPG2500 12-stearoyloxy-stearate***

25

The general reaction route is: 12-hydroxystearic acid → ethyl 12-hydroxystearate → ethyl 12-stearoyloxy-stearate → BuPPG2500 12-stearoyloxy-stearate

Ethyl 12-stearoyloxy-stearate

30 Is obtained in accordance with Example 3.

BuPPG2500 12-stearoyloxy-stearate

10 mmol ethyl 12-stearoyloxy-stearate and 10.5 mmol BuPPG2500 (polypropylene glycol monobutyl ether of average molecular weight 2500 g/mol) are mixed with 500 mg Novozym 435 in a 500 ml round flask with stirring under vacuum (<1 mm Hg) at 75 °C for 100 hours.

The enzyme is removed by filtration to yield BuPPG2500 12-stearoyloxy-stearate.

Example 9*Further examples*

Additional compounds according to the table 2 below were synthesized using the same principal routes, involving the same enzyme, as described above in Examples 1-6.

Table 2.

<u>Compounds</u>	<u>Isolated yield</u>
PEG600 mono-12-hydroxystearate	82%
PEG600 mono-12-acetoxy-stearate	87%
PEG600 mono-12-hexanoyloxy-stearate	73%
PEG600 mono-12-octanoyloxy-stearate	95%
PEG600 mono-12-decanoyloxy-stearate	93%
PEG1500 mono-12-lauroyloxy-stearate	88%
PEG600 mono-12-myristoyloxy-stearate	76%
PEG1500 mono-12-myristoyloxy-stearate	86%
PEG1500 mono-12-palmitoyloxy-stearate	82%
MePEG550 12-hydroxystearate	90%
MePEG1200 12-stearoyloxy-stearate	58%
MePEG2000 12-stearoyloxy-stearate	77%

Isolated yield is calculated based on the fatty acid reactant.

Some of the prepared compounds have been tested as described in Examples 10-12.

5 Example 10.

Haemolytic activity

A static method for evaluating the haemolytic activity of compounds was used. In short, dog blood is processed to an erythrocyte suspension, which is then incubated (40 min at 37°C) with varying concentrations of a surfactant. The incubation is terminated by centrifugation after which the plasma, now containing different amounts of haemoglobin, can be removed and analysed. The results obtained for various MePEG- and PEG mono-12-acyloxy-stearate esters compared with a reference substance (Tween 80), are summarized in tables 3-4.

15

Table 3. Haemolytic activity of monoesters of PEG600, MePEG550 and Tween 80.

PEG600 mono-12-propionyloxy-stearate		PEG600 mono-12-hexanoyloxy-stearate		PEG600 mono-12-octanoyloxy-stearate		MePEG550 12-hydroxystearate		Tween 80	
Conc mM	Hemolysis %	Conc mM	Hemolysis %	Conc mM	Hemolysis %	Conc mM	Hemolysis %	Conc mM	Hemolysis %
0.6	104	0.6	83	0.6	88	1.0	25	1.0	0
		0.7	92	0.7	102	5.0	86	5.0	2
						10	79	10	29

Table 4. Haemolytic activity of monoesters of PEG1500, MePEG1200 and MePEG2000.

PEG1500 mono-12-lauroyloxy-stearate		PEG1500 mono-12-myristoyloxy-stearate		PEG1500 mono-12-palmitoyloxy-stearate		PEG1500 mono-12-stearoyloxy-stearate		PEG1500 mono-12-oleoyloxy-stearate	
Conc mM	Hemolysis %	Conc mM	Hemolysis %	Conc mM	Hemolysis %	Conc mM	Hemolysis %	Conc mM	Hemolysis %
1.0	0	3.1	0	1.0	0	1.0	0	0.8	0
1.5	0	6.2	0	3.0	0	3.1	0	1.9	0
3.0	11	7.7	0	5.0	0	5.1	0	3.7	0
5.0	98	10.8	0	7.0	0	7.2	0	6.0	0
		12.3	0	9.9	0	10.2	0	8.2	0
		15.4	0					11.2	0

MePEG1200 12-palmitoyloxy-stearate		MePEG2000 12-palmitoyloxy-stearate		MePEG1200 12-stearoyloxy-stearate		MePEG2000 12-stearoyloxy-stearate	
Conc mM	Hemolysis %	Conc mM	Hemolysis %	Conc mM	Hemolysis %	Conc mM	Hemolysis %
1	0	1	0	1	0	1	0
5	0	5	0	5	0	5	0
10	0	10	0	10	0	10	0.2

- 5 As shown in Table 3 and Table 4, monoesters with short PEG or MePEG chains (average molecular weight below 1100) caused hemolysis at low concentrations. For example, PEG600 mono-12-propionyloxystearate, PEG600 mono-12-hexanoyloxystearate and PEG600 mono-12-octanoyloxystearate, caused total haemolysis at concentrations below 1 mM. MePEG550 mono-12-hydroxystearate (a compound representing the compounds disclosed in US 6,365,637 B1, Zirnstein et al) caused profound hemolysis at 1 mM and total hemolysis at 5 mM. Furthermore, the commercial product Tween 80 caused haemolysis at concentrations around 5 mM and above. In contrast to the PEG600 monoesters, the monoesters with longer PEG and MePEG chains did not cause any
- 10

haemolysis at all up to 10 mM, except for PEG1500 mono-12-lauroyloxy-stearate which caused total hemolysis already at 5 mM.

5 Example 11

Epithelial cell interaction

Possible interaction with epithelial cells in the form of CACO-2 cells monitored as transepithelial electrical resistance (TEER) was studied for some of the compounds. In
10 short, after an equilibration time of 60 min the buffer solution at the donor side was exchanged with a buffer solution containing a certain concentration of the surfactant and TEER was then measured after 8 hours.

In Table 5 the results for the PEG1500 mono-12-acyloxy-stearate, MePEG1200 12-
15 acyloxy-stearates, MePEG2000 12-acyloxy-stearates and some reference substances tested at different concentrations, are shown. It is evident from Table 5 that even after 8 hours, no change in TEER is observed for monoesters with PEG or MePEG chains longer than 1200 except for PEG1500 mono-12-lauroyloxy-stearate where there was a substantial decrease in TEER indicating some interaction with the epithelial cells. This is in contrast to the short
20 chains of Solutol HS 15, which are not well tolerated.




Table 5. TEER (measured in %) after 8 hours exposure

Compound	Surfactant conc (mM)			
	0.1	1	10	50
Cremophor EL	103	93	50	49
Tween 80	96	99	89	65
Solutol HS15	94	84	37	33
PEG600 mono-12-hexanoyloxy-stearate	89	43	39	37
PEG1500 mono-12-lauroyloxy-stearate	100	96	93	37
PEG1500 mono-12-myristoyloxy-stearate	104	100	104	98
PEG1500 mono-12-palmitoyloxy-stearate	98	105	103	101
PEG1500 mono-12-stearoyloxy-stearate	98	104	103	108
PEG1500 mono-12-oleoyloxy-stearate	105	104	95	108
MePEG1200 12-palmitoyloxy-stearate	-	-	94	103
MePEG1200 12-stearoyloxy-stearate	-	-	101	71
MePEG2000 12-palmitoyloxy-stearate	-	-	96	99
MePEG2000 12-stearoyloxy-stearate	-	-	98	96

- means no measurement performed.

5

Example 12.

Solubilizing capacity

- 10 The solubilization capacity of the compounds of the invention and compounds of prior art for two poorly soluble (i.e. less than sparingly soluble) drug molecules, felodipine and griseofulvin, were tested. The solubility of griseofulvin in water is approximately 23 μM [Karlsson and Söderlind 2000] and the solubility of felodipine in water is 0.0008 mg/ml (2.1 μM , as the $\text{Mw}=384$) [Corswant 1998]. The solubilization capacity for each surfactant
- 15 was determined from the slope of the curve in a graph where the felodipine/griseofulvin concentration was plotted against the surfactant concentration at surfactant concentrations above CMC (critical micellar concentration).

Solubilization capacity for griseofulvin $\alpha_G = \frac{d[G]}{d[S]}$

and for felodipine $\alpha_F = \frac{d[F]}{d[S]}$

where

- 5 [G]=griseofulvin concentration [M]
 [F]=felodipine concentration [M]
 [S]=surfactant concentration [M] (above CMC)

10 The results are summarized in Table 6. The conclusion from these experiments is that the acylated HFA-PEG esters of this invention have better solubilization capacity than the prior art reference, the commercial product Solutol HS 15.

Table 6.

- 15 Solubilization capacity for griseofulvin (α_G) and felodipine (α_F) for the acylated HFA-PEG-esters at 25°C. Solutol HS15 is a commercial product and is used as a reference.

Compound	α_G M/M	α_F M/M
Solutol HS 15	0.015	0.15
PEG1500 mono-12-lauroyloxy-stearate	0.034	0.13
PEG1500 mono-12-myristoyloxy-stearate	0.044	0.31
PEG1500 mono-12-palmitoyloxy-stearate	0.032	0.32
PEG1500 mono-12-stearoyloxy-stearate	0.058	0.29

20 **Example 13.**

Felodipine injectable solution

12 g of PEG1500 mono-12-palmitoyloxy-stearate is dissolved in 988 g saline (0.9 %w/w NaCl in water for injection). 380 mg felodipine is added and the mixture is stirred at room temperature until all felodipine is dissolved. The solution is filled under aseptic conditions into glass vials.

5

Example 14.

Extended release tablets containing 5 mg felodipine.

The tablet are constructed according to the hydrophilic gel matrix principle.

10

7 000 tablets are prepared.

First the solution I (according to below), comprising felodipine (active ingredient), propyl gallate and the surfactant PEG 1500 mono-12-palmitoyloxy-stearate, is made.

15 Secondly, the solution II is prepared.

The powders (III) are mixed in a mixer and moistened with Solution I until homogeneity. The solution II is added and the mixing continued until homogeneity.

20 The obtained granulate is dried in a drying oven.
Thereafter it is milled in a Frewitt oscillating granulator.

25 The granulate obtained is mixed with the powder IV, the lubricant, and mixing is continued for a total of 3 minutes. The mixture is compressed to tablets having an average weight of 227 mg, on a tableting machine, using 9 mm circular concave punches.

		<u>grams</u>
		<u>/7000</u>
	mg/tabl	<u>tablets</u>
Solution I		
Felodipine	5	35
PEG1500 mono-palmitoyloxy-stearate	5	35

30

Propyl gallate	0.06	0.42
Ethanol	30	210
Solution II		
Hydroxypropyl cellulose	10	70
Ethanol	160	1120
Powders III		
Hydroxypropylmethylcellulose 50 cps	100	700
Hydroxypropylmethylcellulose 10000 cps	20	140
Sodium Aluminium Silicate	55	385
Lactose	28	196
Powder IV		
Sodium stearyl fumarate	4.3	30.1

The tablets obtained may be used as cores in e.g. a coating procedure with a pigmented HPMC solution.

5

Example 15.

Extended release tablets containing 5 mg felodipine.

The tablets are constructed according to the hydrophilic gel matrix principle.

10

The tablets are prepared according to Example 14, except that the surfactant used in Example 14, PEG1500 mono-12-palmitoyloxy-stearate is exchanged to MePEG2000 12-stearoyloxy-stearate.

15

Example 16.

Extended release tablets containing 5 mg felodipine.

The tablets are constructed according to the hydrophilic gel matrix principle.

- 5 The tablets are prepared according to Example 14, except that the surfactant used in Example 14, PEG1500 mono-12-palmitoyloxy-stearate is exchanged to MePEG 1200 12-palmitoyloxy-stearate.

Example 17.

- 10 *Extended release tablets containing 5 mg felodipine.*

The tablets are constructed according to the hydrophilic gel matrix principle.

- 15 The tablets are prepared according to Example 14, except that the surfactant used in Example 14, PEG1500 mono-12-palmitoyloxy-stearate is exchanged to PEG1500 mono-12-oleoyloxy-stearate.

20 **Example 18.**

Extended release tablets containing 5 mg felodipine.

The tablets are constructed according to the hydrophilic gel matrix principle.

25

The tablets are prepared according to Example 14, except that the surfactant used in Example 14, PEG1500 mono-12-palmitoyloxy-stearate is exchanged to BuPPG2500 12-stearoyloxy-stearate.

30

Example 19.

Solid dispersion formulation

5 g felodipine is dissolved in 150 g melted MePEG2000 12-stearoyloxy-stearate. The melt (continuously stirred) is filled into hard gelatine capsules. The capsules are cooled under controlled conditions.

5

Example 20.*Emulsion for parenteral use*

- 10 0.5 g felodipine and 1.0 g soybean lecithin are dissolved in 98.5 g soybean oil. 5 g PEG1500 mono-12-palmitoyloxy-stearate is dissolved in 895 g of water for injection. A coarse emulsion is formed by mixing the oil phase with the aqueous phase using an ultraturrax mixer. The droplet size is further decreased by high-pressure homogenisation.

15

Example 21.*Self-emulsifying drug delivery systems (SEDDS)*

- 20 2 g felodipine is dissolved in 300 g of a Miglyol 812/PEG1500 mono-12-palmitoyloxy-stearate mixture (70/30 w/w). The mixture is then filled in soft gelatine capsules.

Example 22.*Suppository formulation*


25

- 5 g felodipine is dissolved in a molten mixture of 175 g PEG1500 mono-12-palmitoyloxy-stearate and 175 g PEG2000 and the mass is cast in appropriate moulds.

30

Claims

1. A compound of the formula (I)



wherein

10 R_1 is H or $\text{C}_1 - \text{C}_4$ alkyl;

R_2 is C_{14} to C_{22} , linear or branched, acyl, alkyl or alkenyl wherein the acyl, alkyl or alkenyl may be optionally further substituted with one or more substituents individually selected from the following; halogen, cyano, carboxy, carbamoyl, carbamoyl($\text{C}_1 - \text{C}_4$)alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, mercapto, nitro, amino, ($\text{C}_1 - \text{C}_4$)alkylamino,

15 phenyl, naphthyl, phenyloxy, naphthyloxy, ($\text{C}_1 - \text{C}_4$)alkylthio, or ($\text{C}_1 - \text{C}_4$)alkylsulfinyl;

R_3 is ethylene, propylene or branched propylene;

x is 2 - 18;

y is 1 - 17;

and the sum of $(x + y)$ is 3 - 19, and

20 z is 25 - 455.

2. A compound according to Claim 1, wherein

R_1 is H or $\text{C}_1 - \text{C}_2$ alkyl.

25 3. A compound according to Claim 1, wherein

x is 2 - 15;

y is 4 - 17;

and the sum of $(x + y)$ is 6 - 19.

30 4. A compound according to Claim 1, wherein

z is 25 - 228.

5. A compound according to Claim 1, wherein

R_1 is H or $C_1 - C_2$ alkyl;

R_2 is C_{14} to C_{22} , linear or branched, acyl, alkyl or alkenyl, wherein the acyl, alkyl or alkenyl may be optionally further substituted with one or more substituents individually selected from the following; halogen, cyano, carboxy, carbamoyl, carbamoyl($C_1 - C_4$)alkyl, fluoromethyl, difluoromethyl, trifluoromethyl, mercapto, nitro, amino, ($C_1 - C_4$)alkylamino, phenyl, naphthyl, phenyloxy, naphthyloxy, ($C_1 - C_4$)alkylthio, or ($C_1 - C_4$)alkylsulfinyl;

R_3 is ethylene, propylene or branched propylene;

x is 2 -15;

y is 4 -17;

and the sum of (x + y) is 6 -19; and

z is 25 - 228.

6. A compound according to any of claims 1-5, wherein

R_1 is H.

7. A compound according to any of claims 1-5, wherein

R_1 is $C_1 - C_2$ alkyl.

8. A compound according to any of claims 1-5, wherein

x is 2 -12;

y is 7 -17;

and the sum of (x + y) is 9 -19.

9. A compound according to any of claims 1-5, wherein

z is 25 - 57.

10. A compound according to claim 5, wherein

R_1 is H or $C_1 - C_2$ alkyl;

R_2 is C_{14} to C_{22} , linear or branched, acyl, alkyl or alkenyl, wherein the acyl, alkyl or alkenyl may be optionally further substituted with one or more substituents individually selected from the following; halogen, cyano, carboxy, carbamoyl, carbamoyl($C_1 - C_4$)alkyl,

fluoromethyl, difluoromethyl, trifluoromethyl, mercapto, nitro, amino, (C₁-C₄)alkylamino, phenyl, naphthyl, phenyloxy, naphthyloxy, (C₁-C₄)alkylthio, or (C₁-C₄)alkylsulfinyl;
R₃ is ethylene, propylene or branched propylene;

x is 2 -12;

5 y is 7 -17;

and the sum of (x + y) is 9 - 19; and

z is 25 - 57.

10

11. A compound according to any of claims 1-10 wherein R₁ is C₁ - C₂ alkyl.

12. A compound according to any of claims 1-10 wherein R₁ is H.

13. A compound according to any of claims 1- 11 wherein R₁ is methyl.

15

14. A formulation comprising a solubilizing compound according to any of claims 1-13 and a compound requiring solubilization.

15. A formulation according to claim 14 where the compound requiring solubilization is a compound having a solubility of less than 33 mg/ml in water.

20

16. A formulation according to claim 14 or 15 comprising a compound according to any of claims 1 - 13, together with a pharmaceutically active ingredient.

17. A formulation according to claim 16 for use as a medicament.

25

18. The use of a compound according to any of claims 1-13, as surfactant in a formulation according to claim 14 or 15, or in a pharmaceutical formulation according to claim 16.

30

19. A process for preparing a polyoxyalkylene glycol (POAG) ester, characterized in that the ester has a poly (oxyalkylene) glycol chain or a C₁ -C₄ alkyl derivatized poly (oxyalkylene) glycol chain having 25 - 455 repeating monomer units and that it utilizes a hydrolytic enzyme catalyzing ester formation with POAG or POAG monoalkyl ether and

the carboxylic acid group of an O-acylated, O-alkylated or O-alkenylated hydroxy fatty acid or C₁ – C₄ alkyl ester without catalyzing any reaction with a bond connecting any acyl, alkyl or alkenyl group to the hydroxy fatty acid or hydroxy fatty acid C₁ – C₄ alkyl ester.

- 5 20. A process according to claim 19 for preparing a compound with formula (I), according
to any of claims 1-13, characterized in that the process utilizes a hydrolytic enzyme
catalyzing ester formation with POAG or POAG monoalkyl ether and the carboxylic acid
group of an O-acylated, O-alkylated or O-alkenylated hydroxy fatty acid or C₁ – C₄ alkyl
ester without catalyzing any reaction with a bond connecting any acyl, alkyl or alkenyl
10 group to the hydroxy fatty acid or hydroxy fatty acid C₁ – C₄ alkyl ester.

21. A process according to claim 20, characterized in that it gives a compound according to
any of claims 1-13, and that it utilizes the hydrolytic enzyme lipase B from *Candida*
antarctica.


15

22. The process according to claim 20, wherein the hydrolytic enzyme is immobilized
lipase B from *Candida antarctica*.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
21

Abstract

New non-ionic surfactants in the form of polyoxyalkylene glycol hydroxy fatty acid derivatives or monoalkylated polyoxyalkylene glycol hydroxy fatty acid derivatives having a polyoxyalkylene / alkyl polyoxyalkylene chain with a chain-length of 25 – 455 repeating
 5 units and a specified substituent on the hydroxy position of the specified hydroxy fatty acid, according to the general formula (I)



wherein for example R_1 is methyl, R_2 is optionally substituted $\text{C}_{14} - \text{C}_{22}$ acyl, alkyl, or
 15 alkenyl, R_3 is ethylene, x is 2-12, y is 7-17, $(x + y)$ is 3-19 and z is 25-57;

advantageously prepared by involving an enzymatic process, formulations comprising them, their use as solubilizers and a process involving a hydrolytic enzyme for preparing them.

